



Quantitative Prediction of Gas-Phase ^{13}C NMR Chemical Shifts

A.A. Auer, J. Gauss

published in

Modern Methods and Algorithms of Quantum Chemistry,
J. Grotendorst (Ed.), John von Neumann Institute for Computing,
Jülich, NIC Series, Vol. 2, ISBN 3-00-005746-3, p. 27, 2000.

© 2000 by John von Neumann Institute for Computing

Permission to make digital or hard copies of portions of this work for personal or classroom use is granted provided that the copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. To copy otherwise requires prior specific permission by the publisher mentioned above.

<http://www.fz-juelich.de/nic-series/>

Quantitative prediction of gas-phase ^{13}C NMR chemical shifts

Alexander A. Auer* and Jürgen Gauss

Universität Mainz
Institut für physikalische Chemie
Jakob-Welder Weg 11
D-55099 Mainz

The accurate prediction of NMR chemical shifts is still a great challenge. While typical errors in SCF ^{13}C chemical shift calculations are about 5-10 ppm, application of correlated methods reduces the errors significantly. Nevertheless it has so far not been possible to predict chemical shifts with an accuracy of about 1 ppm or better and thus to reach quantitative accuracy compared to experimental gas phase NMR data.

In this study we demonstrate that this accuracy can be achieved if :

- chemical shifts are calculated at CCSD(T) level using large basis sets, e.g. of pentuple zeta quality augmented by higher angular momentum functions,
- employed molecular geometries are taken from CCSD(T) optimizations using sufficiently large basis sets such as cc-pVTZ or cc-pVQZ,
- vibrational corrections to chemical shifts are included.

Calculations of ^{13}C chemical shifts for a set of 15 small organic compounds containing different types of bonding environments are presented. A detailed study of the effects of correlation level, basis set convergence and the need of vibrational corrections is carried out in order to reach a maximum deviation of less than 1 ppm.

The results are compared to experimental gas phase ^{13}C NMR data.

* email : auer@mail.uni-mainz.de